

Configuration Interaction calculations of positron binding to $\text{Be}(^3\text{P}^o)$

M.W.J. Bromley

Department of Physics, San Diego State University, San Diego CA 92182, USA

J. Mitroy

Faculty of Technology, Charles Darwin University, Darwin NT 0909, Australia

Abstract

The Configuration Interaction method is applied to investigate the possibility of positron binding to the metastable beryllium ($1s^2 2s 2p\ ^3\text{P}^o$) state. The largest calculation obtained an estimated energy that was unstable by 0.00014 Hartree with respect to the $\text{Ps} + \text{Be}^+(2s)$ lowest dissociation channel. It is likely that positron binding to parent states with non-zero angular momentum is inhibited by centrifugal barriers.

Key words: positron, positron binding, positronic atom, atomic structure, metastable beryllium, configuration interaction

PACS: 36.10.-k, 36.10.Dr, 34.85.+x, 71.60.+z

1 Introduction

In the last several years there has been substantial progress studying the physics of positron binding to atoms. As late as 1997, there was no rigorous evidence for existence of positron-atom bound states [1], although the possibility of positron binding had been often invoked to explain the existence of very large annihilation rates for positrons annihilating in gases [2,3,4,5,6]. Towards the end of 1997, positron binding to atomic lithium was established in two independent calculations [7,8]. Following these initial studies, positron binding to a further 10 elements of the periodic table was demonstrated [9].

Email addresses: mbromley@physics.sdsu.edu (M.W.J. Bromley), jxm107@rsphysse.anu.edu.au (J. Mitroy).

One of the features of these positronic atoms is that all the atoms binding a positron have ionization potentials close to 6.80 eV (the Ps binding energy) and the binding energies are generally larger for the atoms with their ionization energies closest to 6.80 eV.

A related question is whether positrons can form bound states with excited electronic states. There is one excited atomic state known to bind a positron, namely the metastable $\text{He}(^3\text{S}^e)$ state [10,11]. The stability of $e^+\text{He}(^3\text{S}^e)$ state was established with the Stochastic Variational Method (SVM) [12,13,7] following an initial inconclusive study by Drachman *et al* [14].

The answer to this question is relevant to positron-atom collision physics [15], since such states may manifest themselves as Feshbach resonances in the elastic or excitation cross sections. There has been a suggestion that the rich resonant structures prominent in electron-atom scattering [16] are largely absent from the positron-atom spectrum [17,9,15].

This work investigates the possibility of positron binding to the low-lying triplet odd state of neutral beryllium, i.e. the $\text{Be}(1s^2 2s 2p\ ^3\text{P}^o)$ system. This state has been identified as one of the most likely excited states to bind a positron [18,9]. This conjecture has largely been based upon the fact that its ionization potential (IP) of 6.598 eV [19] is quite close to the Ps binding energy. This IP is between that of magnesium and calcium, both of which bind a positron with a binding energy of ≈ 0.4 eV [9,20,21]. Furthermore, the dipole polarisability of $\text{Be}(^3\text{P}^o)$ is $\alpha_d = 39.02\ a_0^3$ [22], greater than that of ground state Be ($\alpha_d = 37.7\ a_0^3$ [23]).

This metastable state warrants investigation for another reason; the parent atom has an angular momentum greater than zero with the loosely bound $2p$ electron carrying the angular momentum. The quantitative impact that the angular momenta has on the ability to bind a positron is largely unknown. Angular momentum coupling considerations suggests that the existence of a repulsive centrifugal barrier will act to inhibit positron binding by making it less energetically favorable to form a Ps-cluster. Either, the electron-positron pair will be in a relative p -state with a smaller energy gain than the $\text{Ps}(1s)$ state, or the total angular momentum of the Ps cluster relative to the nucleus will be greater than zero.

2 Details of the Calculation

Since the ionization energy of the $\text{Be}(2s 2p\ ^3\text{P}^o)$ system is just less than 6.80 eV, the threshold for a stable positron complex is that for the $\text{Be}^+(2s) + \text{Ps}(1s)$ dissociation channel. This energy was -0.9192086 Hartree for the present

model potential. An energy level diagram showing the ground and low-lying metastable states of Be, and some negative ion and positronic ion energies are shown in Figure 1.

The CI method as applied to positron-atomic systems with two valence electrons and a positron has been discussed previously [24,25,20], but a short description is worthwhile. The model Hamiltonian is initially based on a Hartree-Fock (HF) wave function for the neutral atom ground state. One- and two-body semi-empirical polarization potentials are added to the potential field of the HF core and the parameters of the core-polarization potentials defined by reference to the spectrum of Be^+ [26].

All calculations were done in the frozen-core approximation. The general effective Hamiltonian for the system with N_e valence electrons and a positron was

$$H = -\frac{1}{2}\nabla_0^2 - \sum_{i=1}^{N_e} \frac{1}{2}\nabla_i^2 + \sum_{i<j}^{N_e} \frac{1}{r_{ij}} + \sum_{i=1}^{N_e} (V_{dir}(\mathbf{r}_i) + V_{exc}(\mathbf{r}_i) + V_{p1}(\mathbf{r}_i)) \\ - \sum_{i=1}^{N_e} \frac{1}{r_{i0}} - V_{dir}(\mathbf{r}_0) + V_{p1}(\mathbf{r}_0) - \sum_{i<j}^{N_e} V_{p2}(\mathbf{r}_i, \mathbf{r}_j) + \sum_{i=1}^{N_e} V_{p2}(\mathbf{r}_i, \mathbf{r}_0) . \quad (1)$$

The direct potential (V_{dir}) represents the interaction with the HF $1s^2$ electron core. The direct part of the core potential is attractive for electrons and repulsive for the positron. The exchange potential (V_{exc}) between the valence electrons and the HF core was computed without approximation.

The one-body polarization potential (V_{p1}) was a semi-empirical polarization potential derived from an analysis of the Be^+ spectrum. It has the functional form

$$V_{p1}(r) = -\frac{\alpha_d g^2(r)}{2r^4} . \quad (2)$$

The factor α_d is the static dipole polarizability of the core and $g^2(r)$ is a cutoff function designed to make the polarization potential finite at the origin. The same cutoff function has been adopted for both the positron and electrons. In this work, $g^2(r)$ was defined to be

$$g^2(r) = 1 - \exp(-r^6/\rho^6) , \quad (3)$$

where ρ is an adjustable cutoff parameter. The core dipole polarizability was set to $0.0523 a_0^3$ while ρ was set to $0.95 a_0$ [25]. The two-body polarization potential (V_{p2}) is defined as

$$V_{p2}(\mathbf{r}_i, \mathbf{r}_j) = \frac{\alpha_d}{r_i^3 r_j^3} (\mathbf{r}_i \cdot \mathbf{r}_j) g(r_i) g(r_j) . \quad (4)$$

This model potential gives a $\text{Be}(2s2p\ ^3P^o)$ binding energy of 0.91147 Hartree with respect to the Be^{2+} threshold. The experimental binding energy with respect to this threshold is 0.91888 Hartree [27].

The CI basis was constructed by letting the two electrons and the positron form all the possible total angular momentum $L_T = 1$ configurations, with the two electrons in a spin-triplet state, subject to the selection rules, namely

$$\max(\ell_0, \ell_1, \ell_2) \leq L_{\max} \quad (5)$$

$$\min(\ell_1, \ell_2) \leq L_{\text{int}} \quad (6)$$

$$(-1)^{(\ell_0+\ell_1+\ell_2)} = -1 \quad (7)$$

In these rules ℓ_0 , ℓ_1 and ℓ_2 are respectively the orbital angular momenta of the positron and the two electrons.

Our two-electron-positron calculations with non-zero total angular momentum were first validated against the previous $L_T = 1$ and 2 PsH calculations of Tachikawa [28]. Using their Gaussian-type orbitals we reproduced their reported energy and annihilation rates. Note that the PsH states with $L_T = 1$ and 2 are unbound [29,28,30].

For the $e^+\text{Be}(^3P^o)$ calculations, the Hamiltonian was diagonalized in a CI basis constructed from a very large number of single particle orbitals, including orbitals up to $\ell = 12$. There was a minimum of 12 radial basis functions for each ℓ . The largest calculation was performed with $L_{\max} = 12$ and $L_{\text{int}} = 3$ and gave a CI basis dimension of 498750. The resulting Hamiltonian matrix was diagonalized with the Davidson algorithm [31], and a total of 1376 iterations were required for the largest calculation.

3 Results

Various $e^+\text{Be}(^3P^o)$ expectation values, such as energy and mean distance of the electron and positron from the nucleus are given in Table 1. The 2γ annihilation rate [9] for annihilation with the core and valence electrons are denoted Γ_c and Γ_v respectively. The calculations shown in Table 1 have a minimum of 12 radial basis functions for each ℓ with $L_{\max} = 12$ and $L_{\text{int}} = 3$. The largest calculation remains unbound by -0.0021904 Hartree.

To demonstrate that $L_{\text{int}} = 3$ is sufficient for this system, also shown in Table 1 is the result of a $L_{\max} = 12$ and $L_{\text{int}} = 2$ calculation. The $L_{\max}$ parameter needs to be large since it determines the extent to which electron-positron correlations are incorporated into the wavefunction. However, L_{int} is largely concerned with electron-electron correlations and, for example, setting $L_{\text{int}} = 3$

for the PsH and $e^+\text{Be}$ ground states recovered 99.4% and 98.5% respectively of the Ps and positron binding energies for a given L_{max} [25].

The main problem afflicting CI calculations of positron-atom interactions is the slow convergence of the expectation values with L_{max} [32,33,9,34]. One way to determine the $L_{\text{max}} \rightarrow \infty$ expectation values is to assume that the successive increments, ΔX_L , to any expectation value $\langle X \rangle$ scale as $1/L^p$ as L increases [32]. However, arguments based on 2nd-order perturbation theory suggests that the energy increments scale asymptotically as $1/(L + \frac{1}{2})^{p_E} \approx 1/(L + \frac{1}{2})^4$ [35,36]. While existing CI calculations are consistent with this idea [25,37,34,38], the actual size of the exponent, p_E is significantly smaller than 4 at $L_{\text{max}} \approx 10$. This can be seen in Table 1, where the power law factor p_E relating successive $E(L_{\text{max}} - 2)$, $E(L_{\text{max}} - 1)$, and $E(L_{\text{max}})$ are tabulated. Applying a simple power law extrapolation with $p_E = 2.078$ leads to a prediction of binding ($E = -0.9207021$), but not much credence should be placed in this since fixing p_E at its $L_{\text{max}} = 12$ value of 2.078 will result in an extrapolation that will overestimate the contribution of the orbitals with $L > L_{\text{max}}$ [39,34].

An improved method, borrowing from existing work in atomic structure physics [40,41,21], is to estimate the $L_{\text{max}} \rightarrow \infty$ limit for the energy by assuming that

$$\Delta E_L \approx \frac{A_E}{(L + \frac{1}{2})^4} + \frac{B_E}{(L + \frac{1}{2})^5} + \frac{C_E}{(L + \frac{1}{2})^6}. \quad (8)$$

The factors A_E , B_E and C_E are determined from four calculations at successively larger values of L_{max} . Applying eq. (8) to the data in Table 1 at $L_{\text{max}} = 12$ one finds $A_E = -34.601$, $B_E = 477.71$ and $C_E = -1829.3$, with $E = -0.919070$ Hartree as our present best energy estimate. This is 0.0001390 Hartree higher the threshold for binding at -0.9192086 Hartree.

Usage of an inverse power series restricted to the first two terms of eq. (8) results in $A_E = -21.875$ and $B_E = 172.29$, and an $e^+\text{Be}(^3\text{P}^o)$ energy of $E = -0.9188267$, which is also unbound. As has been seen in other CI calculations of positron-atom systems, even when the three-term and the two-term coefficients are significantly different, their extrapolated energies can still lie relatively close together [34].

There are two aspects where convergence can be incomplete. Besides the size of L_{max} , the number of LTOs for each value of ℓ still needs to be further increased. This is best seen in Figure 2 which depicts the energies as a function of L_{max} . Two sets of $e^+\text{Be}(^3\text{P}^o)$ data are shown, the first shows the $e^+\text{Be}(^3\text{P}^o)$ energies from Table 1 which used 12 LTOs, and their extrapolated estimates using eq. (8). The second set shows the results from initial exploratory calculations also with $L_{\text{max}} = 12$ and $L_{\text{int}} = 3$, but only with a minimum of 8 LTOs per ℓ . Even a cursory glance at Figure 2 suggests that the CI prognosis for variationally establishing the existence of a bound state is not promising. Also

shown in Figure 2 are the $\text{Be}(^3\text{P}^o)$ energies using the underlying $e^+\text{Be}(^3\text{P}^o)$ basis without the positron (the 12 LTOs basis).

Even though the valence annihilation rate from the largest calculation is only $0.348 \times 10^9 \text{ sec}^{-1}$, the slow convergence of Γ_v with L_{max} suggests the presence of a well defined Ps cluster. Fitting the four largest Γ_v calculations in Table 1 to the three-term form

$$\Delta\Gamma_L \approx \frac{A_\Gamma}{(L + \frac{1}{2})^2} + \frac{B_\Gamma}{(L + \frac{1}{2})^3} + \frac{C_\Gamma}{(L + \frac{1}{2})^4}, \quad (9)$$

results in an extrapolated annihilation rate of $\Gamma_v = 0.805 \times 10^9 \text{ sec}^{-1}$. However, this value should be regarded as only notional. After all, if this system is indeed unbound, Γ_v should approach $\Gamma_{\text{Ps}} = 2.008 \times 10^9 \text{ sec}^{-1}$ in the limit of an infinite radial basis [42].

It is also noticed that $\langle r_p \rangle$ in Table 1 steadily decreases as L_{max} increases. This also should not be taken as an indicator of binding since the behavior of $\langle r_p \rangle$ with L_{max} is not straightforward. For example, the PsH $\langle r_p \rangle$ decreases when L_{max} changes from 0 to 3, but then starts increasing as L_{max} increases from 3 to 9 [25,34]. Indeed, this is a trend noticed in CI calculations of positronic atoms with $IP < 6.8\text{eV}$ (eg. $e^+\text{Li}$ [43], $e^+\text{Ca}$ and $e^+\text{Sr}$ [20,38]), which have shown an initial $\langle r_p \rangle$ decrease to a minimum, which then increases as $L_{\text{max}} \rightarrow \infty$. The behavior of $\langle r_p \rangle$ with L_{max} for an unbound system is unknown.

4 Conclusions

A large-scale CI calculation of the $e^+\text{Be}(^3\text{P}^o)$ ground state has been performed. The largest calculation gave an energy that was 0.00219 Hartree above the threshold for binding. Using an extrapolation method to estimate the contribution from the higher-partial waves gave an energy that was 0.00014 Hartree above threshold. By its very nature the present calculation is unable to give definitive proof that the $e^+\text{Be}(^3\text{P}^o)$ system does not have a bound state. While a variational calculation of the present kind can be used to give proof of binding, it cannot be used to give proof of non-binding. A converged calculation of low energy $\text{Be}^+\text{-Ps}(1s)$ scattering in the $^4\text{P}^o$ channel would be needed to establish lack of binding.

What can be inferred from the present calculation is that a $e^+\text{Be}(^3\text{P}^o)$ bound state (assuming one existed) would have a structure similar to positronic lithium, i.e. it would consist of a Ps-like object weakly bound to the $\text{Be}^+(2s)$ core. The size of the CI expansion required to establish binding would be enormous. For example, it is necessary to include orbitals with $L_{\text{max}} \approx 30$ to get an energy that is lower than the $\text{Ps} + \text{Li}^+$ threshold [43]. Another conclusion

is that the existence of centrifugal barriers does seem to inhibit the binding of positrons to parent atoms with non-zero angular momentum.

Acknowledgments

The authors would like to thank Shane Caple of CDU for computer support. Some preliminary calculations were performed on computing facilities made possible by the Research Corporation. One author (MB) would like to thank Dr. Brett Esry for discussions on three-body systems with non-zero angular momentum. The authors would also like to thank Dr. Masanori Tachikawa for access to unpublished data on positronic systems with non-zero angular momentum.

References

- [1] D. M. Schrader, Bound States of Positrons with Atoms and Molecules: Theory, Nucl. Instrum. Methods Phys. Res. B 143 (1998) 209.
- [2] D. A. L. Paul, L. Saint-Pierre, Rapid Annihilations of Positrons in Polyatomic Gases, Phys. Rev. Lett. 11 (1963) 493.
- [3] H. C. Khare, P. R. Wallace, G. G. Bach, A. Chodos, Annihilation of positrons in liquid helium and the positron-helium bound state, Can. J. Phys. 42 (1964) 1522.
- [4] V. I. Goldanskii, Y. S. Sayasov, On the resonance annihilation of positrons in collisions of neutral atoms or molecules, Phys. Lett. 13 (1964) 300.
- [5] J. D. McNutt, D. A. Johnson, V. B. Summerour, Annihilations of Positrons in Methane, Phys. Rev. B 4 (1971) 36.
- [6] C. M. Surko, A. Passner, M. Leventhal, F. J. Wysocki, Bound states of positrons and large molecules, Phys. Rev. Lett. 61 (1988) 1831.
- [7] G. G. Ryzhikh, J. Mitroy, Positronic lithium, an electronically stable Li-e^+ ground state, Phys. Rev. Lett. 79 (1997) 4124.
- [8] K. Strasburger, H. Chojnacki, Quantum chemical study of simple positronic systems using explicitly correlated Gaussian functions - PsH and PsLi^+ , J. Chem. Phys. 108 (1998) 3218.
- [9] J. Mitroy, M. W. J. Bromley, G. G. Ryzhikh, Positron and positronium binding to atoms and ions, J. Phys. B 35 (2002) R81.
- [10] G. G. Ryzhikh, J. Mitroy, A metastable state of positronic helium, J. Phys. B 31 (1998) 3465.

- [11] J. Mitroy, Expectation values of the $e^+\text{He}(^3\text{S}^e)$ system, *Phys. Rev. A* 72 (2005) 032503.
- [12] K. Varga, Y. Suzuki, Precise solution of few-body problems with the stochastic variational method on a correlated Gaussian basis, *Phys. Rev. C* 52 (1995) 2885.
- [13] Y. Suzuki, K. Varga, *Approach to Quantum-Mechanical Few-Body Problems*, 172, Springer, New York, 1998.
- [14] R. J. Drachman, Y. K. Ho, S. K. Houston, Positron attachment to helium in the ^3S state, *J. Phys. B* 9 (1976) L199.
- [15] C. M. Surko, G. F. Gribakin, S. J. Buckman, Low-energy positron interactions with atoms and molecules, *J. Phys. B* 38 (2005) R57.
- [16] S. J. Buckman, C. W. Clark, Atomic negative-ion resonances, *Rev. Mod. Phys.* 66 (1994) 539.
- [17] J. P. Sullivan, S. J. Gilbert, S. J. Buckman, C. M. Surko, Searches for resonances in the scattering of low-energy positrons from atoms and molecules, *J. Phys. B* 34 (2001) L467.
- [18] D. M. Schrader, Antimatter Compounds, in: C. M. Surko, F. A. Gianturco (Eds.), *New Directions in Antimatter Physics and Chemistry*, Kluwer Academic Publishers, The Netherlands, 2001, p. 263.
- [19] S. Bashkin, J. O. Stoner, *Atomic Energy Levels and Grotrian Diagrams*, North Holland, Amsterdam, 1975.
- [20] M. W. J. Bromley, J. Mitroy, Configuration Interaction calculations of positron binding to group II elements, *Phys. Rev. A* 65 (2002) 062505.
- [21] M. W. J. Bromley, J. Mitroy, Convergence of partial wave expansion of the He ground state (2006) in preparation.
- [22] J. Mitroy, M. W. J. Bromley, Properties of the triplet metastable states of alkaline-earth-metal atoms, *Phys. Rev. A* 70 (2004) 052503.
- [23] J. Mitroy, M. W. J. Bromley, Semi-empirical calculation of van der Waals coefficients for alkali-metal and alkaline-earth-metal atoms, *Phys. Rev. A* 68 (2003) 052714.
- [24] M. W. J. Bromley, J. Mitroy, G. G. Ryzhikh, Configuration interaction calculations of positronic atoms and ions, *Nucl. Instrum. Methods Phys. Res. B* 171 (2000) 47.
- [25] M. W. J. Bromley, J. Mitroy, Configuration Interaction calculations of PsH and $e^+\text{Be}$, *Phys. Rev. A* 65 (2002) 012505.
- [26] D. W. Norcross, M. J. Seaton, Energy levels for Be I calculated using a model potential and cores approximations, *J. Phys. B* 9 (1976) 2983.
- [27] NIST Atomic Spectra Database Version 3.0 (2005).
URL [http://physics.nist.gov/cgi-bin/AtData/main\\$_\\$asd](http://physics.nist.gov/cgi-bin/AtData/main$_$asd)

- [28] M. Tachikawa, Simultaneous optimization of Gaussian type function exponents for electron and positron with full-CI wavefunction - application to ground and excited states of positronic compounds with multi-component molecular orbital approach, *Chem. Phys. Lett.* 350 (2001) 269.
- [29] D. Bressanini, M. Mella, G. Morosi, Stability and positron annihilation of positronium hydride $L = 0, 1, 2$ states: A quantum Monte Carlo study, *Phys. Rev. A* 57 (1998) 1678.
- [30] S. L. Saito, Multireference configuration interaction calculations of some low-lying states of positronium hydride, *J. Chem. Phys.* 118 (2003) 1714.
- [31] A. Stathopoulos, C. Froese Fischer, A davidson program for finding a few selected extreme eigenpairs of a large, sparse, real, symmetric matrix, *Comput. Phys. Commun.* 79 (1994) 268.
- [32] J. Mitroy, G. G. Ryzhikh, Configuration interaction calculation of the Cu^+ ground state, *J. Phys. B* 32 (1999) 2831.
- [33] V. A. Dzuba, V. V. Flambaum, G. F. Gribakin, C. Harabati, Calculation of the positron bound state with the copper atom, *Phys. Rev. A* 60 (1999) 3641.
- [34] J. Mitroy, M. W. J. Bromley, Convergence of CI single center calculations of positron-atom interactions (2006) in preparation.
- [35] C. Schwartz, Importance of Angular Correlations between Atomic Electrons, *Phys. Rev.* 126 (1962) 1015.
- [36] G. F. Gribakin, J. Ludlow, Convergence of partial-wave expansions for energies scattering amplitude and positron annihilation rates, *J. Phys. B* 35 (2002) 339.
- [37] M. W. J. Bromley, J. Mitroy, Variational calculation of positron-atom scattering using configuration-interaction-type wave functions, *Phys. Rev. A* 67 (2003) 062709.
- [38] M. W. J. Bromley, J. Mitroy, Large dimension Configuration Interaction calculations of positron binding to the group II atoms, *Phys. Rev. A* 72 (2005) in prep.
- [39] M. W. J. Bromley, J. Mitroy, Configuration Interaction calculations of positron binding to Zn and Cd, *Phys. Rev. A* 65 (2002) 062506.
- [40] D. P. Carroll, H. J. Silverstone, R. P. Metzger, Piecewise polynomial configuration interaction natural orbital study of $1s^2$ helium, *J. Chem. Phys.* 71 (1979) 4142.
- [41] R. N. Hill, Rates of convergence and error estimation formulas for the Rayleigh-Ritz variational method, *J. Chem. Phys.* 83 (1985) 1173.
- [42] J. Mitroy, M. W. J. Bromley, Comment on: Multi-reference configuration interaction calculations for positronium halides [*J. Chem. Phys.* **122**, 054302 (2005)], *J. Chem. Phys.* 123 (2005) 017101.

- [43] M. W. J. Bromley, J. Mitroy, Positron and positronium interactions with Cu, Phys. Rev. A 66 (2002) 062504.
- [44] T. Andersen, H. H. Andersen, P. Balling, P. Kristensen, V. V. Petrunin, Structure and dynamics of the negative alkaline-earth ions, J. Phys. B 30 (1997) 3317.
- [45] J. Mitroy, G. G. Ryzhikh, Improved description of LiPs, e^+ Be, NaPs and e^+ Mg, J. Phys. B 34 (2001) 2001.




Fig. 1. The energy levels of electron and positron binding to neutral beryllium (in units of Hartree relative to the Be^{2+} threshold). The experimental atomic binding energies are taken from the NIST compilation [27], the metastable negative ion experimental binding energies are from [44], and the theoretical e^+ Be binding energy is taken from a frozen-core SVM calculation [45]. The threshold for e^+ binding to $\text{Be}(^3\text{P}^o)$ is denoted as $\text{Be}^+ + \text{Ps}$.

Table 1. Results of CI calculations for $e^+\text{Be}(^3\text{P}^o)$ for a series of L_{max} , with fixed $L_{\text{int}} = 3$ (the row denoted 12* was computed with $L_{\text{int}} = 2$). The total number of electron and positron orbitals are identical and denoted by N . The 3-body energy of the $e^+\text{-Be}(^3\text{P}^o)$ system, relative to the energy of the Be^{2+} core, is denoted by E (in Hartree). The threshold for binding is -0.9192086 Hartree, and ε gives binding energy (in Hartree) against dissociation into $\text{Ps} + \text{Be}^+(2s)$. The mean electron-nucleus distance $\langle r_e \rangle$, the mean positron-nucleus distance $\langle r_p \rangle$, and the mean electron-positron separation $\langle r_{ep}^2 \rangle$ are given in units of a_0 . The Γ_v and Γ_c columns give the valence and core annihilation rates respectively (in units of 10^9 sec^{-1}). The p_E column gives the power-series exponents from the $L_{\text{max}}, L_{\text{max}}-1, L_{\text{max}}-2$ energies. The results in the row ∞ used eq. (8) and eq. (9) to estimate the $L_{\text{max}} \rightarrow \infty$ correction.

L_{max}	N	N_{CI}	E	ε	$\langle r_e \rangle$	$\langle r_p \rangle$	$\langle r_{ep}^2 \rangle$	Γ_c	Γ_v	p_E
1	32	11325	-0.9049669	-0.0142417	2.70529	31.8236	1159.699	0.000020	0.000521	
2	45	32580	-0.9105271	-0.0086815	2.70601	27.4961	920.4328	0.000087	0.004302	
3	57	65694	-0.9112555	-0.0079531	2.72410	21.5600	619.5881	0.000284	0.020168	6.0406
4	69	108570	-0.9120770	-0.0071316	2.76099	16.3218	378.4203	0.000571	0.053933	-0.479
5	81	153510	-0.9129784	-0.0062302	2.80833	13.2441	250.9955	0.000816	0.097421	-0.462
6	93	201606	-0.9138427	-0.0053659	2.85800	11.6135	189.6602	0.000977	0.141967	0.2515
7	105	250350	-0.9146093	-0.0045993	2.90693	10.6898	157.1977	0.001076	0.184291	0.8378
8	117	300030	-0.9152671	-0.0039415	2.95389	10.1253	138.1528	0.001133	0.223404	1.2235
9	129	349710	-0.9158238	-0.0033848	2.99833	9.75778	125.9701	0.001166	0.259160	1.5001
10	141	399390	-0.9162926	-0.0029160	3.03992	9.50665	117.6386	0.001183	0.291698	1.7182
11	153	449070	-0.9166867	-0.0025219	3.07841	9.32966	111.6870	0.001191	0.321221	1.9055
12	165	498750	-0.9170182	-0.0021904	3.11361	9.20197	107.2937	0.001193	0.347951	2.0775
12*	165	334248	-0.9169903	-0.0022183	3.11398	9.20391	107.3440	0.001192	0.347880	
∞			-0.9190697	-0.0001390					0.805027	




Fig. 2. The energy (in units of Hartree) of e^+ -Be($^3P^o$) (crosses) and extrapolated energies (diamonds) as a function of L_{max} for two different calculations (with a minimum of 8 and 12 LTOs per ℓ respectively). The Be($^3P^o$) two-electron energy is shown by the squares for the 12 LTOs basis. The threshold for binding at $E(\text{Be}^+)+E(\text{Ps})$ is shown as solid line.